

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:		ModHomCo (Pty) Ltd						Version:	02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions	
1	Imported and Exported	Berberis-Homaccord	Liquid	30 mL	Each 30 mL cont.: Berberis vulgaris L. D4 0,1 mL; Berberis vulgaris L. D10 0,1 mL; Berberis vulgaris L. D30 0,1 mL; Berberis vulgaris L. D200 0,1 mL; Citrillus colocynthus (L.) Schrad D4 0,1 mL; Citrillus colocynthus (L.) Schrad D10 0,1 mL; Citrillus colocynthus (L.) Schrad D30 0,1 mL; Citrillus colocynthus (L.) Schrad D200 0,1 mL; Veratrum album L. D4 0,1 mL; Veratrum album L. D10 0,1 mL; Veratrum album L. D30 0,1 mL; Veratrum album L. D200 0,1 mL.	The usual dose is: Adults: Take 10 drops 3 times daily. Hold in mouth 10 to 15 minutes before swallowing. In acute disorders, take 10 drops every 15 minutes for 8 doses. Children and infants: Half the adult dosage. The drops may be taken in a little water. It is recommended not to eat 15 minutes before or after taking the medication.	U5548 D540530	Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in inflammation or irritation (with or without concretions) in the region of the urogenital tract and the biliary duct.	At or below 25 ° C	
2	Imported and Exported	*[S3] Cerebrum compositum NM	Injectable (D5 only)	10 x 2,2 mL	2,2 mL contains: Acidum Phosphoricum D10 22 mg; Aconitum napellus L. D6 22 mg; Aesculus hippocastanum L. D4 22 mg; Physeter macrocephalus L. (Ambr. grisea) D10 22 mg; Anamirta cocculus (L.) Wight & Arn. D4 22 mg; Arnica montana L. D28 22 mg; Sus scrofa ferus domesticus (Cerebrum suis) D8 22 mg; Cinchona pubescens Vahl (Cinchona succirubra Pav.), Cinchona calisaya Wedd., Cinchona ledgeriana Moens ex Trimen, (Cinchona pubescens mother tincture) D4 22 mg; Conium maculatum L. D4 22 mg; Sus scrofa ferus domesticus (Embryo totalis suis) D10 22 mg; Gelsemium sempervirens (L.) J.St.-Hil D4 22 mg; Sus scrofa ferus domesticus (Hepar suis) D10 22 mg; Hyoscyamus niger L. D6 22 mg; Kalium bichromicum D8 22 mg; Kalium phosphoricum D6 22 mg; Magnesium phosphoricum D10 22 mg; Sus scrofa ferus domesticus (Placenta totalis suis) D10 22 mg; Ruta graveolens L. D4 22 mg; Selenium D10 22 mg; Semecarpus anacardium L. D6 22 mg; Strychnos ignatii P.J. Bergius D8 22 mg; Sulfur D10 22 mg; Thuja occidentalis L. D6 22 mg; Manganum phosphoricum D8 "aquaos".	The usual dose is: In general, 1 ampoule 1-3 times weekly I.m., s.c., i.d., or if necessary i.v.	U5642 D540452	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in the stimulation of the defence mechanism in dysontogenesis in children, legasthenia, vegetative dystonia, depression, arteriosclerosis, after commotio cerebri, after encephalitis, in neuralgia, states of agitation, amyotrophic lateral sclerosis, multiple sclerosis, Parkinson's disease, deficiency of memory, nervous exhaustion and other geriatric indications.	At or below 25 ° C	
3	Imported and Exported	*[S3] Coenzyme compositum	Injectable (D5 only)	10 x 2,2 mL	2,2 mL cont.: Coenzymum A D8 22 µl; Acidum ascorbicum D6 22 µl; Thiaminum hydrochloricum D6 22 µl; Natrium riboflavinum phosphoricum D6 22 µl; Pyridoxinum hydrochloricum D6 22 µl; Nicotinamidum D6 22 µl; Acidum cis-aconiticum D8 22 µl; Acidum citricum D8 22 µl; Acidum fumaricum D8 22 µl; Acidum alphaketoglutaricum D8 22 µl; Acidum DL-malicum D8 22 µl; Acidum succinicum D8 22 µl; Barium oxalsuccinicum D10 22 µl; Natrium diethylloxalacetikum D6 22 µl; Natrium pyruvicum D8 22 µl; Cysteinum D6 22 µl; Pulsatilla pratensis D6 22 µl; Hepar sulfuris D10 22 µl; Sulfur D10 22 µl; Adenosinum triphosphoricum D10 22 µl; Nadidum D8 22 µl; Manganum phosphoricum D6 22 µl; Magnesium oroticum D6 22 µl; Cerium oxalicum D8 22 µl; Acidum alpha-liponicum D6 22 µl; Beta vulgaris conditiva D4 22 µl.	The usual dose is: Adults: In general, 1 ampoule 1-3 times weekly I.m., s.c., i.c. Infants (birth to 4 years old): Inject 0,7 mL Children ages 4-8: Inject 1 mL and ages 8-12, inject 1,3 mL. Discard any unused solution.	U5643 D540451	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for stimulating blocked enzymatic systems in degenerative diseases as well as in defective enzymatic functions (cellular phases).	At or below 25 ° C	
4	Imported and Exported	Cralonin	Liquid	30 mL	30 mL cont.: Crataegus species (Crataegus oxyacantha) Ø HAB 21,0 g, including: Crataegus laevigata (Poir.) or Crataegus monogyna Jacq.; Spigelia anthelmia L. D3 0,3 g; Potassium carbonate (Kalium carbonicum) 0,3 g.	The usual dose is: Typical dosage for long term therapy: Adults: Take 10-20 drops 3 times daily. Hold in mouth 10-15 seconds before swallowing. In acute disorders, take 10 drops every 15 minutes (Caution: the patient should not take the medication in this dosage over lengthy periods of time without control by a physician). Shake before use. The drops may be taken in a little water. It is recommended not to eat or drink 15 minutes before or after taking this medication.		Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in senile heart, consequences of damage to the cardiac musculature, nervous cardiac disorders, piercing cardiac pain, and cardiac pain of other nature (e.g. anginose complaints).	At or below 25 ° C	

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:		ModHomCo (Pty) Ltd			Version:			02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions
5	Imported and Exported	*[S3] Discus compositum	Injectable (DS only)	10 x 2,2 mL	2,2 mL cont.: Acidum ascorbicum (ascorbic acid) D6 22,0 mg; Acidum picricum (picric acid) D6 22,0 mg; Acidum silicium D6 22,0 mg; Acidum thiocticum (thioctic acid) D8 22,0 mg; Aesculus hippocastanum L. D6 22,0 mg; Ammonium chloratum D8 22,0 mg; Argentum metallicum (silver powder) D10 22,0 mg; Berberis vulgaris L. D4 22,0 mg; Calcium phosphoricum D10 22,0 mg; Cartilago suis (Sus scrofa ferus domesticus) D8 22,0 mg; Cnicifraga racemosa (L.) Nutt. D4 22,0 mg; Cinchona pubescens Vahl D4 22,0 mg; Citrus aurantium D4 22,0 mg; Citrullus colocynthis (L.) Schrad. D4 22,0 mg; Coenzyme A D10 22,0 mg; Cuprum aceticum (copper (II) acetate monohydrate) D6 22,0 mg; Discus intervertebralis suis (Sus scrofa ferus domesticus) D8 22,0 mg; Embryo totalis suis (Sus scrofa ferus domesticus) D10; Funiculus umbilicalis suis (Sus scrofa ferus domesticus) D10 22,0 mg; Glandula suprarenalis suis (Sus scrofa ferus domesticus) D10 22,0 mg; Hydragrynum oxydatum rubrum (red mercury (II) oxide) D10 22,0 mg; Kalium carbonicum (potassium carbonate) D6 22,0 mg; Ledum palustre L. D4 22,0 mg; Medorrhinum nosode D18 22,0 mg; Medulla ossis suis (Sus scrofa ferus domesticus) D10 22,0 mg; Natrium diethyloxalacetum (sodium diethyloxalacetate) D6 22,0 mg; Natrium riboflavinum phosphoricum (riboflavin sodium phosphate) D6 22,0 mg; Nicotinamidum (nicotinamide) D6 22,0 mg; Pseudogalphalum obtusifolium (L.) Hillard & B.L. Burt D3 22,0 mg; Pulsatilla pratensis (L.) Mill. D6 22,0 mg; Pyridoxinum hydrochloricum D6 22,0 mg; Ranunculus bulbosus L. D4 22,0 mg; Scilla corniculata (Claviceps purpurea (Fries) Tulane) D6 22,0 mg; Sepia officinalis L. D10 22,0 mg; Sulfur D28 22,0 mg; Thiaminum hydrochloricum (thiamine hydrochloride) D6 22,0 mg;	The usual dose is: In general, 1 ampoule 1-3 times weekly i.m., s.c., i.c. (neural).	U5649 D540464	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use as stimulation of the endogenic defence reactions and influencing of the homotoxic terrain in osteochondrosis, disorders of the joints (chronic arthritis and arthrosis), particularly for euralgic/rheumatic disorders in the region of the vertebral column.	At or below 25 °C
6	Imported and Exported	*[S3] Echinacea compositum S	Injectable (DS only)	10 x 2,2 mL	2,2 mL cont.: Echinacea angustifolia D3 22 µL; Aconitum napellus L. D3 22 µL; Sanguinaria canadensis D4 22 µL; Sulfur D8 22 µL; Baptisia tinctoria D4 22 µL; Lachesis mutio D10 22 µL; Bryonia cretica D6 22 µL; Eupatorium perfoliatum D6 22 µL; Pulsatilla pratensis D8 22 µL; Mercurius sublimatus corrosivus D8 22 µL; Thuja occidentalis D8 22 µL; Influenzinum-Nosode D13 22 µL; Phosphorus D8 22 µL; *[4] Cortisonum aceticum [4] D13 22 µL; Streptococcus haemolyticus-Nosode D18 22 µL; Staphylococcus-Nosode D18 22 µL; Phytolacca americana D6 22 µL; Pyrogenium-Nosode D198 22 µL; Zincum metallicum D10 22 µL; Gelsemium sempervirens L. J.St.-Hill D6 22 µL; Hepar sulfuris D10 22 µL; Rhus toxicodendron D4 22 µL; Arnica montana L. D4 22 µL; Arsenicum album D8 22 µL; Argentum nitricum D8 22 µL; Euphorbium D6 22 µL.	The usual dose is: Adults: For the general stimulation of the defensive system: 1 ampoule should be administered 1-3 times weekly i.m., s.c., i.d. or, if necessary, i.v. The intramuscular or intravenous injection and dose must be adapted to the individual case and the reaction of the patient. Infants (birth to 4 years old): Inject 0,7 ml. Children aged 4-8 years: Inject 1 ml and ages 8-12 years, inject 1,3 ml. Discard any unused solution.	U5650 D510189	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use to stimulate the body's intrinsic defence in cases of fever and inflammation, postvaccinal encephalitis, influenza, angina, furuncles, abscesses, phlegmons, focal toxicosis, gingivitis, stomatitis, sinusitis, gastroenteritis, enterocolitis, cystitis, pyelitis, colitis, leucorrhea, adnexitis, glomerulonephritis, fistular suppurations, osteomyelitis, otitis media, chronic cerebral abscess, meningitis, anthrax, carbuncles, mononucleosis, skin diseases, mastitis.	At or below 25 °C
7	Imported and Exported	Engystol	Tablet	50	1 tabl. cont.: The active substances are: Vincetoxicum hirundinaria D6 75 mg; Vincetoxicum hirundinaria D10 75 mg; Vincetoxicum hirundinaria D30 75 mg; Sulfur D4 37,5 mg; Sulfur D10 37,5 mg. The other ingredient is: Magnesium stearate. Each tablet contains approx. 300 mg of lactose monohydrate.	The usual dose is: Adults and children 12 years and older: In general, 1 tablet 3 x daily. Children 2-6 years: In general, 1 tablet 2 x daily. The tablets should be allowed to dissolve slowly in the mouth. Infants up to 2 years: In general, 1 tablet 1 x daily. For children and infants, it is recommended to crush the tablet and administer it dissolved in a little water. Do not eat or drink 15 minutes before or after taking the medication.	D501098	Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed to activate the non-specific defensive mechanism, particularly in the case of influenza and questionable virus diseases.	At or below 25 °C
8	Imported and Exported	*[S3] Engystol N	Injectable (DS only)	10 x 1,1 mL	1,1 mL cont.: The active substances are: Vincetoxicum hirundinaria D6 6,6 µL; Vincetoxicum hirundinaria D10 6,6 µL; Vincetoxicum hirundinaria D30 6,6 µL; Sulfur D4 3,3 µL; Sulfur D10 3,3 µL.	The usual dose is: Adult: In acute disorders 1 ampoule daily, otherwise 3-1 times weekly i.m., s.c., i.c., i.v. Infants (birth to 4 years old): Inject 0,3 ml. Children aged 4-8: Inject 0,5 ml and ages 8-12, inject 0,7 ml. Discard any unused solution.	U5588 D501099	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed to activate the non-specific defensive mechanism, particularly in the case of influenza and questionable virus diseases.	At or below 25 °C

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:	ModHomCo (Pty) Ltd					Version:		02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions
9	Imported and Exported	Gripp-Heel	Tablet	50	1 tablet cont.: The active substances are: Aconitum napellus D4 120 mg; Bryonia cretica D4 60 mg; Lachesis mutus D12 60 mg; Eupatorium perfoliatum D3 30 mg; Phosphorus D5 30 mg. The other ingredient is: Magnesium stearate. 1 tablet contains approx. 300 mg lactose.	The usual dose is: Adults and children over 3 years of age: Dissolve 1 tablet in the mouth 3-5 times daily. In acute disorders, 1 tablet every 15 minutes for 8 doses. Infants: Half the adult dose. Crush the tablet and administer it dissolved in a little water or milk. It is recommended not to eat or drink 15 minutes before or after taking the medication.	U5496 D540598	Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in influenza and influenza infections and for the stimulation of the endogenous defence system in other feverish infectious diseases.	At or below 25 ° C
10	Imported and Exported	*[S3] Gripp-Heel	Injectable (DS only)	10 x 1,1 mL	1,1 mL cont.: The active substances are: Aconitum napellus D3 4,4 µl; Eupatorium perfoliatum D2 1,1 µl; Phosphorus D4 1,1 µl; Lachesis mutus D11 2,2 µl; Bryonia cretica D3 2,2 µl.	The usual dose is: Adults: In acute disorders 1 ampoule daily, otherwise 1 ampoule 1-3 times weekly i.m., s.c., i.c., i.v.. Infants (birth to 4 years old): Inject 0.3 ml. Children ages 4-8: Inject 0.5 ml and ages 8-12, inject 0.7 ml. Discard any unused solution.	U5591 D540532	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in cases of influenza and influenza infections.	At or below 25 ° C
11	Imported and Exported	*[S3] Hepar compositum	Injectable (DS only)	10 x 2,2 mL	2,2 mL cont.: The active substances are: Hepar suis D8 22 µl; Cyanocobalaminum D4 22 µl; Duodenum suis D10 22 µl; Thymus suis D10 22 µl; Colon suis D10 22 µl; Vesica fellea suis D10 22 µl; Pankreas suis D10 22 µl; China D4 22 µl; Lycopodium clavatum D4 22 µl; Chelidonium majus D4 22 µl; Carduus marianus D3 22 µl; Histaminum D10 22 µl; Sulfur D13 22 µl; Avena sativa D6 22 µl; Fel tauri D8 22 µl; Natrium diethylsacacetum D10 22 µl; Acidum alphaketoglutaricum D10 22 µl; Acidum DL-malicum D10 22 µl; Acidum fumaricum D10 22 µl; Acidum alpha-liponicum D8 22 µl; Acidum oroticum D6 22 µl; Cholesterinum D10 22 µl; Calcium carbonicum Hahnemanni D28 22 µl; Taraxacum officinale D4 22 µl; Cynara scolymus D6 22 µl; Veratrum album D4 22 µl.	The usual dose is: In general, 1 ampoule 1-3 times weekly i.m., s.c., i.c., or, if necessary, i.v.	U5654 D540658	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use as stimulation of the detoxification function of the liver in acute and chronic diseases of the liver/gall, e.g. cholangitis and cholecystitis; in functional disorders of the liver of toxic origin as cause or result of other diseases, in hypercholesteremia, rythematodes, toxæanthema, dermatosis and dermatitis, neurodermitis and for the stimulation of the liver cell functioning.	At or below 25 ° C
12	Imported and Exported	*[S3] Lymphomyosot N	Injectable (DS only)	10 x 1,1 mL	1,1 mL cont.: Araneus diadematus D6 0,555 mg; Calcium phosphoricum (calcium hydrogen phosphate dehydrate) D12 0,55 mg; Equisetum hyemale L. D4 0,55 mg; Ferrum iodatum (iron (II) iodide on lactose D12 1,10 mg; Fumaria officinalis L. D4 0,55 mg; Gentiana lutea L. D5 0,55 mg; Geranium robertianum L. D4 1,10 mg; *[4] Levothyroxium (levothyroxine free acid) D12 0,55 mg; Myosotis arvensis (L.) Hill. D3 0,55 mg; Nasturtium officinale R.Br. D4 1,10 mg; Natrium sulfuricum (anhydrous sodium sulphate) D4 0,55 mg; Pinus sylvestris L. D4 0,55 mg; Scophularia nodosa L. 0,55 mg; Smilax ornata Lem. D6 0,55 mg; Teucrium scorodonia L. 0,55 mg; Veronica officinalis L. D3 0,55 mg.	The usual dose is: In acute disorders, inject 1 ampoule daily, otherwise 1 ampoule 1-3 times weekly i.m., s.c., i.v..	U5625	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in lymphatism (tendency to hypertrophy of the lymphatic organs, tendency to formation of oedemas and to susceptibility to infection); glandular swelling; tonsillar hypertrophy and chronic tonsillitis.	At or below 25 ° C

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:		ModHomCo (Pty) Ltd						Version:	02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions	
13	Imported and Exported	Lymphomyosot N	Liquid	30 mL	30 mL cont.: Araneus diadematus D6 0,05 mL; Calcium phosphoricum (calcium hydrogen phosphate dehydrate) D12 0,05 mL; Equisetum hyemale L. D4 0,05 mL; Ferrum iodatum (Iron (II) iodide on lactose) D12 0,10 mL; Fumaria officinalis L. D4 0,05 mL; Gentiana lutea L. D4 0,10 mL; Geranium robertianum L. D4 0,10 mL; *[4] Levothyroxinum (levothyroxine free acid) D12 0,05 mL Myosotis arvensis (L.) Hill. D3 0,05 mL; Nasturtium officinale R.Br. D4 0,10 mL; Natrium sulfuricum (anhydrous sodium sulphate) D4 0,05 mL; Pinus sylvestris L. D4 0,05 mL; Scrophularia nodosa L. D3 0,05 mL; Smilax ornata Lem. D60,05 mL; Teucrium scorodonia L. D3 0,05 mL; Veronica officinalis L. D3 0,05 mL. Contains 35 % v/v ethyl alcohol.	In general, 15-20 drops 3 times daily. The drops may be taken in a little water and should be held in the mouth for 10-15 seconds before swallowing. It is recommended not to eat or drink 15 minutes before or after taking the medication. Replace cap tightly after use.	U5530 D540661	Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in lymphatism (tendency to hypertrophy of the lymphatic organs, tendency to formation of oedemas and to susceptibility to infection); glandular swelling; tonsillar hypertrophy and chronic tonsillitis.	At or below 25 ° C	
14	Imported and Exported	*[S3] Mucosa compositum	Injectable (D5 only)	10 x 2,2 mL	2,2 mL cont.: The active substances are: Mucosa nasalis suis D8 22 µl; Mucosa oris suis D8 22 µl; Mucosa pulmonis suis D8 22 µl; Mucosa oculi suis D8 22 µl; Mucosa vesicae felleae suis D8 22 µl; Mucosa vesicae urinariae suis D8 22 µl; Mucosa pylori suis D8 22 µl; Mucosa duodeni suis D8 22 µl; Mucosa oesophagi suis D8 22 µl; Mucosa jejuni suis D8 22 µl; Mucosa ileitis suis D8 22 µl; Mucosa coli suis D8 22 µl; Mucosa recti suis D8 22 µl; Mucosa ductus choledochi suis D8 22 µl; Ventriculus suis D8 22 µl; Pancreas suis D10 22 µl; Argentum nitricum D8 22 µl; Atropa belladonna D10 22 µl; Oxalis acetosella D6 22 µl; Semecarpus anacardium D6 22 µl; Phosphorus D8 22 µl; Lachesis mutus D10 22 µl; Ipecacuanha D8 22 µl; Nux vomica D13 22 µl; Veratrum album D4 22 µl; Pulsatilla pratensis D6 22 µl; Kreosotum D10 22 µl; Sulfur D8 22 µl; Natrium diethylacetikum D8 22 µl; Bacterium coli- Nosode D28 22 µl; Marsdenia cundurango D6 22 µl; Kalium bichromicum D8 22 µl; Hydrastis canadensis D4 22 µl; Mandragora e radice siccato D10 22 µl; Momordica balsamina D6 22 µl;	The usual dose is: In acute disorders 1 ampoule daily, otherwise 1 ampoule 1-3 times weekly i.m., s.c., i.c., or if necessary i.v..	U5658	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use as stimulation of the bodily defences in diseases of the mucous membranes and catarrhs of various types and localisation, e.g. in the alimentary tract with or without ulcers, in the region of the upper and lower parts of the respiratory tract, the urinary passages and the conjunctiva.	At or below 25 ° C	
15	Imported and Exported	Nervoheel N	Tablet	50	Each tablet contains: The active substances are: Acidum phosphoricum D4 60 mg; Strychnos ignatii D4 60 mg; Sepia officinalis D4 60 mg; Kalium bromatum D4 30 mg; Zincum isovalerianicum D4 30 mg. The other ingredients are: Magnesium stearate, Lactose monohydrate (300 mg).	The usual dose is: In general, dissolve one tablet in the mouth 3 times daily. It is recommended not to eat or drink 15 minutes before or after taking the medication.	U5503	Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in patients suffering from general exhaustion and nervous complaints.		

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:		ModHomCo (Pty) Ltd					Version:		02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions	
16	Imported and Exported	Neurexan	Tablet	50	Each tablet contains: The active substances are: Passiflora incarnata D2 0,6 mg; Avena sativa D2 0,6 mg; Coffea arabica D12 0,6 mg; Zincum isovalerianicum D4 0,6 mg. The other ingredient is: Magnesium stearate. Each tablet contains 300 mg lactose.	The usual dose is: Adults: Acute: In general, 1 tablet to be dissolved slowly in the mouth every half to one hour. Do not exceed 12 tablets in one day for long-term therapy. Chronic: 1 tablet 1-3 times daily. Insomnia: Adults (12 years and older): In general, 1 tablet 3 times during the evening. If insomnia persists, 1 tablet every 15 minutes, up to 3 more times. Children (6 to 11 years): In general, 1 tablet twice during the evening. If insomnia persists, 1 tablet every 15 minutes, up to 2 more times. Children (1 to 5 years): In general, 1/2 tablet 3 times during the evening. If insomnia persists 1/2 tablet every 15 minutes, up to 3 more times. Babies (0 to 12 months): In general, 1/2 tablet twice during the evening. If insomnia persists, 1/2 tablet every 15 minutes, up to 2 more times. Dissolve tablet in a teaspoon of water before administering to a small child.	D552010	Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is indicated for nervous restlessness, insomnia and sleeping problems.	At or below 25 ° C	
17	Imported and Exported	Nux vomica-Homaccord	Liquid	30 mL	30 mL cont.: The active substances are: Nux vomica D2 0,06 mL; Nux vomica D10 0,06 mL; Nux vomica D15 0,06 mL; Nux vomica D30 0,06 mL; Nux vomica D200 0,06 mL; Nux vomica D1000 0,06 mL; Bryonia cretica D2 0,06 mL; Bryonia cretica D6 0,2 mL; Bryonia cretica D10 0,06 mL; Bryonia cretica D15 0,06 mL; Bryonia cretica D30 0,06 mL; Bryonia cretica D200 0,06 mL; Bryonia cretica D1000 0,06 mL; Lycopodium clavatum D3 0,09 mL; Lycopodium clavatum D10 0,09 mL; Lycopodium clavatum D30 0,09 mL; Lycopodium clavatum D200 0,09 mL; Lycopodium clavatum D1000 0,09 mL; Citrullus colocynthis D3 0,09 mL; Citrullus colocynthis D10 0,09 mL; Citrullus colocynthis D30 0,09 mL; Citrullus colocynthis D200 0,09 mL. The other ingredients are: Ethanol 96 %, Purified water. Contains 35 % v/v ethyl alcohol.	The usual dose is: Adults: Take 10 drops 3 times daily. Hold in mouth 10-15 seconds before swallowing. In acute disorders, take 10 drops every 15 minutes for 8 doses. Long-term administration of this preparation (for several months) must be supervised by a therapist. Children and infants: Half the adult dose. The drops may be taken in a little water. It is recommended not to eat or drink 15 minutes before or after taking the medication.	U5567 D552011	Discipline specific	Homeopathy	Oral	Low	This medicine is prepared in accordance with homeopathic principles and is proposed for use in functional disorders in the gastrointestinal and hepatic region; meteorism; disorders after consumption of alcohol, coffee or nicotine.	At or below 25 ° C	

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:		ModHomCo (Pty) Ltd						Version:		02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities		Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions	
18	Imported and Exported	*[S3] Ovarium compositum	Injectable (DS only)	10 x 2,2 mL	2,2 mL cont.: The active substances are: Ovarium suis D8 22 µl; Placenta suis D10 22 µl; Uterus suis D10 22 µl; Salpinx suis D10 22 µl; Hypophysis suis D13 22 µl; Cypripedium calceolus var. pubescens D6 22 µl; Lilium tigrinum D4 22 µl; Pulsatilla pratensis D18 22 µl; Aquilegia vulgaris D4 22 µl; Sepia officinalis D10 22 µl; Lachesis mutus D10 22 µl; Apisinum D8 22 µl; Kreosotum D8 22 µl; Bovista D6 22 µl; Ipecacuanha D6 22 µl; Mercurius solubilis Hahnemanni D10 22 µl; Hydrastis canadensis D4 22 µl; Acidum cis-aconiticum D10 22 µl; Magnesium phosphoricum D10 22 µl.		The usual dose is: In acute disorders: 1 ampoule daily, otherwise 1 ampoule 1-3 times weekly i.m., s.c., i.c., or if necessary, i.v.	U5659	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use as stimulation of the glandular and defensive functions, as well as those of the connective tissue, in dysmenorrhoea, endometritis, metritis, parametritis, enuresis (in young girls), in the climacteric, hyperemesis, insufficiency of the anterior lobe of the pituitary gland in females, craurosis vulvae, mastodynia, osteomalacia, menorrhagia, as well as in various disturbances of metabolism, including those arising in geriatrics.	At or below 25 ° C	
19	Imported and Exported	Reneel	Tablet	50	1 tablet cont.: The active substances are: Berberis vulgaris D2 15 mg; Acidum nitricum D4 30 mg; Cantharis D5 30 mg; Plumbum acetatum D6 30 mg; Pareira brava D3 30 mg; Sabal serrulatum D2 30 mg; Causticum Hahnemanni D4 60 mg; Aluminium oxydatum D12 75 mg. The other ingredient is: Magnesium stearate. Contains lactose monohydrate.		The usual dose is: Adults and children over 3 years of age: Dissolve 1 tablet in the mouth 3 times daily. In acute disorders, 1 tablet every 15 minutes for 8 doses. Infants: Half the adult dose. Crush the tablet and administer it dissolved in a little water or milk. It is recommended not to eat or drink 15 minutes before or after taking the medication.	U5508	Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in inflammatory diseases in the region of the urinary passages, with and without lithiasis.	At or below 25 ° C	
20	Imported and Exported	*[S3] Spascupreel	Injectable (DS only)	10 x 1,1 mL	1,1 mL cont.: The active substances are: Citrullus colocynthis D4 1,1 µl; Ammonium bromatum D4 1,1 µl; Atropinum sulfuricum D6 1,1 µl; Veratrum album D6 1,1 µl; Magnesium phosphoricum D6 1,1 µl; Gelsemium sempervirens D6 1,1 µl; Passiflora incarnata D2 0,55 µl; Agaricus D4 0,55 µl; Chamomilla recutita D3 0,55 µl; Cuprum sulfuricum D6 0,55 µl; Aconitum napellus D6 2,2 µl.		The usual dose is: In acute disorders, 1 ampoule several times daily, otherwise 1 ampoule 1-3 times weekly i.m., s.c., i.c., i.v.	U5630 D540531	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in case of spasms of the smooth musculature of the hollow organs (stomach, intestine, gall bladder, uterus, urinary tract); spasticity of the striated musculature (myogelosis, hardening of the muscles).	At or below 25 ° C	
21	Imported and Exported	Spascupreel	Tablet	50	1 tablet cont.: The active substances are: Citrullus colocynthis D4 30 mg; Ammonium bromatum D4 30 mg; Atropinum sulfuricum D6 30 mg; Veratrum album D6 30 mg; Magnesium phosphoricum D6 30 mg; Gelsemium sempervirens D6 30 mg; Passiflora incarnata D2 15 mg; Agaricus D4 15 mg; Chamomilla recutita D3 15 mg; Cuprum sulfuricum D6 15 mg; Aconitum napellus D6 60 mg. The other ingredient is: Magnesium stearate. One tablet contains approx. 300 mg lactose.		The usual dose is: Adults and children over 3 years of age: Dissolve 1 tablet in the mouth 3 times daily. In acute disorders, 1 tablet every 15 minutes for 8 doses. Infants: Half the adult dose. Crush the tablet and administer it dissolved in a little water or milk. It is recommended not to eat or drink 15 minutes before or after taking the medication.	U5510 D510040		Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in spasms in the organs of the smooth musculature (stomach, intestine, gall bladder, uterus, urinary tract); spasticity of the striated musculature (myogelosis, hardening of the muscles).	At or below 25 ° C	

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:	ModHomCo (Pty) Ltd						Version:	02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions
22	Imported and Exported	*[S3] Testis compositum N	Injectable (DS only)	10 x 2,2 mL	Each 2,2 mL ampoule contains: The active substances are: Testis suis D4 22 mg; Embryo suis D8 22 mg; Glandula suprarenalis suis D13 22 mg; Kalium picrinicum D6 22 mg; Panax ginseng D4 22 mg; Turnera diffusa D8 22 mg; Caladium seguinum D6 22 mg; Cor suis D8 22 mg; *[4] Cortisonum aceticum D13 22 mg; Vitex agnus-castus D6 22 mg; Selenium D10 22 mg; Lytta vesicatoria D8 22 mg; Conium maculatum D28 22 mg; Lycopodium clavatum D28 22 mg; Phosphorus D8 22 mg; Diencephalon suis D10 22 mg; Magnesium phosphoricum D10 22 mg; Ferrum phosphoricum D10 22 mg; Zincum metallicum D10 22 mg; Acidum ascorbicum D6 22 mg; Manganum phosphoricum D8 22 mg.	The usual dose is: In general, 1 ampoule 1-3 times weekly i.m., s.c., i.d..	U5667 D540659	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in disorders of the male sexual function such as impotence, in the stimulation of the male organs in dystrophia adipsogenitalis, ejaculatio praecox (premature ejaculation), impotentia virilis, nocturnal enuresis, dystrophia musculorum progressiva, exhaustion, induration penis plastica, osteomalacia, diabetes mellitus and in other various metabolic disturbances, damaged liver. Stimulation of the regressive vicariation.	At or below 25 ° C
23	Imported and Exported	*[S3] Thyreidea compositum	Injectable (DS only)	10 x 2,2 mL	2,2 mL cont.: The active substances are: Glandula thyreidea suis D8 22 µl; Thymus suis D10 22 µl; Corpus pineale suis D8 22 µl; Splen suis D10 22 µl; Medulla ossis suis D10 22 µl; Funiculus umbilicalis suis D10 22 µl; Hepar suis D10 22 µl; Galium aparine D4 22 µl; Sedum acre D6 22 µl; Sempervivum tectorum D6 22 µl; Conium maculatum D4 22 µl; Spongia D8 22 µl; Acidum sarcolacticum D3 22 µl; Fucus vesiculosus D6 22 µl; Calcium fluoratum D10 22 µl; Colchicum autumnale D4 22 µl; Viscum album D3 22 µl; *[4] Cortisonum aceticum D28 22 µl; Pulsatilla pratensis D8 22 µl; Sulfur D10 22 µl; Natrium diethylloxalacetikum D8 22 µl; Acidum fumaricum D8 22 µl Acidum DL-malicum D8 22 µl; Acidum alpha- ketoglutaricum D8 22 µl; Adenosinum triphosphoricum D8 22 µl.	The usual dose is: In general, 1 ampoule 1-3 times weekly i.m., s.c., i.c., or if necessary i.v.	U5667 D540659	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in case of regulation of the defence system in connective tissue, thyroid and other glandular dysfunctions.	At or below 25 ° C
24	Imported and Exported	*[S3] Tonico-Injeel	Injectable (DS only)	10 x 1,1 mL	1,1 mL cont.: The active substances are: Aurum metallicum D10 1,65 µl; Aurum metallicum D30 1,65 µl; Aurum metallicum D200 1,65 µl; Chininum arsenicosum D10 1,1 µl; Chininum arsenicosum D30 1,1 µl; Helonias dioica D10 1,1 µl; Helonias dioica D30 1,1 µl; Hypericum perforatum D10 1,1 µl; Hypericum perforatum D30 1,1 µl; Selenium D10 1,1 µl; Selenium D30 1,1 µl; Kalium phosphoricum D10 2,2 µl; Kalium phosphoricum D30 2,2 µl; Nux vomica D10 2,75 µl; Nux vomica D30 2,75 µl.	The usual dose is: In acute disorders 1 ampoule daily, otherwise 1 ampoule 1-3 times weekly i.m., s.c., i.c., i.v.	U5632	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in case of states of exhaustion of physical as well as psychic nature. Mental development deficiencies among children.	At or below 25 ° C

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:		ModHomCo (Pty) Ltd						Version:		02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions		
25	Imported and Exported	*[S3] Traumeel S	Injectable (D5 only)	10 x 2,2 mL	Each 2,2 mL (= 2,2 g) ampoule contains: The active substances are: Arnica montana D2 2,2 µl; Calendula officinalis D2 2,2 µl; Chamomilla recutita D3 2,2 µl; Symphytum officinale D6 2,2 µl; Achillea millefolium D3 2,2 µl; Atropa belladonna D2 2,2 µl; Aconitum napellus D2 1,32 µl; Bellis perennis D2 1,1 µl; Hypericum perforatum D2 0,66 µl; Echinacea angustifolia D2 0,55 µl; Echinacea purpurea D2 0,55 µl; Hamamelis virginiana D1 0,22 µl; Mercurius solubilis Hahnemannii D6 1,1 mg; Hepar sulfuris D6 2,2 µl.	The usual dose is: Adults and Children above 6 years: In acute disorders daily, otherwise 1-3 times weekly 1-2 ampoules i.m., s.c., i.e. (in the case of wheals), i.v., intra-articular or periarticular. Infants and Children to 6 years: Half the adult dose.	U5633	Discipline specific	Homeopathy	Injection	High	Traumeel® S is indicated in injuries such as sprains, dislocations, contusions, effusions of blood and effusions into a joint, fractures; post-operative and post-traumatic oedema and swelling of the soft tissues; inflammatory processes and degenerative processes associated with inflammation on the various organs and tissues, including, in particular, on the support and mobility apparatus (tendovaginitis, styloiditis, epicondylitis, bursitis, scapulohumeral periarthritis); arthrosis of the hip, knee and small joints; commotio cerebri acuta.	At or below 25 ° C		
26	Imported and Exported	Traumeel S	Gels	50 g	50 g gel contains: The active substances are: Arnica montana L. D3 0,75 g; Calendula officinalis Ø 0,225 g; Hamamelis virginiana Ø 0,225 g; Echinacea angustifolia Ø 0,075 g; Echinacea purpurea Ø 0,075 g; Chamomilla recutita Ø 0,075 g; Symphytum officinale D4 0,05 g; Bellis perennis Ø 0,05 g; Hypericum perforatum D6 0,045 g; Achillea millefolium Ø 0,045 g; Aconitum napellus L. D1 0,025 g; Atropa belladonna D1 0,025 g; Mercurius solubilis Hahnemannii D6 0,020 g; Hepar sulfuris D6 0,0125 g.	For external use only. The usual dose is: Apply to the affected parts 1-2 times daily, or more often if necessary. A dressing may be applied. Do not use on open wounds.	None	Discipline specific	Homeopathy	Cutaneous	High	For the temporary relief of arthritic pain, of symptoms associated with injuries (i.e. sports injuries, sprains, bruises) and inflammations of all kinds (i.e. tennis elbow).	At or below 25 ° C		
27	Imported and Exported	Traumeel S	Tablet	1 x 50 1 x 250	1 tablet contains: The active substances are: Arnica montana D2 15 mg; Calendula officinalis D2 15 mg; Hamamelis virginiana D2 15 mg; Achillea millefolium D3 15 mg; Atropa belladonna D4 75 mg; Aconitum napellus D3 30 mg; Mercurius solubilis Hahnemannii D8 30 mg; Hepar sulfuris D8 30 mg; Chamomilla recutita D3 24 mg; Symphytum officinale D8 24 mg; Bellis perennis D2 6 mg; Echinacea angustifolia D2 6 mg; Echinacea purpurea D2 6 mg; Hypericum perforatum D2 3 mg.	The usual dose is: Adults and children over 3 years of age: Dissolve 1 tablet in the mouth 3 times daily. Infants: Half the adult dose.	U5514	Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in injuries such as sprains, dislocations, contusions, effusions of blood and effusions into a joint, fractures; post-operative and post-traumatic oedema and swelling of the soft tissues; inflammatory processes and degenerative processes associated with inflammation on the various organs and tissues, including, in particular, on the support and mobility apparatus (tendovaginitis, styloiditis, epicondylitis, bursitis, scapulohumeral periarthritis); arthrosis of the hip, knee and small joints; commotio cerebri acuta.	At or below 25 ° C		
28	Imported and Exported	Traumeel S	Liquid	1 x 30 mL	Each 30 mL contain: The active substances are: Arnica montana L. D2 1,5 mL; Calendula officinalis D2 1,5 mL; Hamamelis virginiana D2 1,5 mL; Achillea millefolium D3 1,5 mL; Atropa belladonna D4 7,5 mL; Aconitum napellus L. D3 3,0 mL; Mercurius solubilis Hahnemannii D8 3,0 mL; Hepar sulfuris D8 3,0 mL; Chamomilla recutita D3 2,4 mL; Symphytum officinale D8 2,4 mL; Bellis perennis D2 0,6 mL; Echinacea angustifolia D2 0,6 mL; Echinacea purpurea D2 0,6 mL; Hypericum perforatum D2 0,3 mL. The other ingredient is: Purified water. Contains 35 % v/v ethyl alcohol.	The usual dose is: Adults and Children above 11 years: 10 drops orally 3 times daily; for swelling of the soft tissues 30 drops 3 times daily. Hold in mouth 10-15 seconds before swallowing. Infants and Children to 11 years: Due to its alcohol content (35 vol.-% v/v ethyl alcohol), a medical practitioner should be consulted before using Traumeel® S for children below 12 years. For best results, Traumeel® S should be administered on an empty stomach.	U5540	Discipline specific	Homeopathy	Oral	High	Traumeel® S is indicated in injuries such as sprains, dislocations, contusions, effusions of blood and effusions into a joint, fractures; post-operative and post-traumatic oedema and swelling of the soft tissues; inflammatory processes and degenerative processes associated with inflammation on the various organs and tissues, including, in particular, on the support and mobility apparatus (tendovaginitis, styloiditis, epicondylitis, bursitis, scapulohumeral periarthritis); arthrosis of the hip, knee and small joints; commotio cerebri acuta.	At or below 25 ° C		

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:		ModHomCo (Pty) Ltd						Version:	02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions	
29	Imported and Exported	Traumeel S	Other	1 x 50 g	Each 50 g cont.: The active substances are: Arnica montana L. D3 0.75 g; Calendula officinalis Ø 0.225 g; Hamamelis virginiana Ø 0.225 g; Echinacea angustifolia Ø 0.075 g; Echinacea purpurea Ø 0.075 g; Chamomilla recutita Ø 0.075 g; Symphytum officinale D4 0.05 g; Bellis perennis Ø 0.05 g; Hypericum perforatum D6 0.045 g; Achillea millefolium Ø 0.045 g; Aconitum napellus L. D1 0.025 g; Atropa belladonna D1 0.025 g; Mercurius solubilis Hahnemanni D6 0.02 g; Hepar sulfuris D6 0.0125 g. The other ingredients are: Cetostearyl alcohol (type A) emulsifying; Ethanol (96 per cent); Paraffin, liquid; Paraffin, white soft; Water, purified. Preservative: 13,8 % v/v ethyl alcohol.	The usual dose is: Adults, children and infants: apply ointment generously to the affected area 2 to 3 times daily. Traumeel® S should be rubbed gently into the skin. Traumeel® S may be applied using mild compression bandaging and/or occlusive bandaging. Sufficient ointment should be applied to cover the affected area, but should not be applied over large areas. Avoid contact with eyes, mucosae, open wounds or broken skin.	U5684	Discipline specific	Homeopathy	Cutaneous	High	Traumeel® S is indicated in injuries of all kinds (sports, accidents) such as sprains, dislocations, contusions, effusions of blood and effusions into a joint, fractures, etc; inflammatory processes and degenerative processes associated with inflammation on the various organs and tissues (e.g. periodontitis, suppuration of the gingival pockets, periodontosis), including, in particular, on the support and mobility apparatus (tendovaginitis, bursitis, scapulohumeral periarthritis); arthrosis of the hip, knee and small joints.	At or below 25 ° C	
30	Imported and Exported	*[S3] Ubichinon compositum	Injectable (DS only)	10 x 2,2 mL	2,2 mL cont.: The active substances are: Ubichinonum D10 22 µL; Acidum ascorbicum D6 22 µL; Thiaminum hydrochloricum D6 22 µL; Natrium riboflavinum phosphoricum D6 22 µL; Pyridoxinum hydrochloricum D6 22 µL; Nicotinamidum D6 22 µL; Vaccinium myrtillus D4 22 µL; Colchicum autumnale D4 L. 22 µL; Podophyllum peltatum L. D4 22 µL; Conium maculatum L. D4 22 µL; Hydrastis canadensis D4 22 µL; Acidum sarcolacticum D6 22 µL; Hydrochinonum D8 22 µL; Acidum alpha-liponicum D8 22 µL; Sulfur D8 22 µL; Manganum phosphoricum D8 22 µL; Natrium diethyloxalacetikum D8 22 µL; Trichinoylum D10 22 µL; Anthrachinonum D10 22 µL; Naphthochinonum D10 22 µL; para-Benzochinonum D10 22 µL; Adenosinum triphosphoricum D10 22 µL; Coenzymum A D10 22 µL; Galium aparine D6 22 µL; Acidum acetylosalicylicum D10 22 µL; Histaminum D10 22 µL; Nadidum D10 22 µL; Magnesium gluconicum D10 22 µL.	The usual dose is: Adults: In general, 1 ampoule 1-3 times weekly i.m., s.c., i.c. Infants (birth to 4 years old): Inject 0,7 ml. Children ages 4-8: Inject 1 ml and ages 8-12, inject 1,3 ml.	U5671 D510191	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for stimulating the defensive mechanisms against toxins in order to reactivate the blocked enzymatic systems in defective enzymatic functions and degenerative diseases (cellular phases).	At or below 25 ° C	
31	Imported and Exported	Vertigoheel	Tablet	1 x 50	1 tablet cont.: The active substances are: Anamirta cocculus D4 210 mg; Conium maculatum D3 30 mg; Ambra grisea D6 30 mg; Petroleum rectificatum D8 30 mg. The other ingredient is: Magnesium stearate. Contains lactose monohydrate.	The usual dose is: Adults and children over 3 years of age: Dissolve 3 tablets in the mouth 3 times daily. Infants: Half the adult dose. Crush the tablet and administer it dissolved in a little water or milk. It is recommended not to eat or drink 15 minutes before or after taking the medication.	U5515	Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in dizziness of various origins (particularly arising from arteriosclerosis).	At or below 25 ° C	
32	Imported and Exported	Viburcol N	Other	1 x12	Each suppository contains: The active substances are: Chamomilla recutita D1 1,100 mg; Atropa belladonna D2 1,100 mg; Plantago major D3 1,100 mg; Pulsatilla pratensis D2 2,200 mg; Calcium carbonicum Hahnemanni D8 4,400 mg. The other ingredient is: Hard fat.	The usual dose is: Remove suppository from plastic foil. In acute disorders, insert one suppository rectally and repeat on several occasions; after alleviation of symptoms, insert one suppository 2-3 times daily. For infants (from the first day of life) up to six months of age, do not use more than one suppository twice daily.	U5677	Discipline specific	Homeopathy	Rectal	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in infants and children in conditions of restlessness with or without fever; symptomatic treatment of common infections.	At or below 25 ° C	

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:		ModHomCo (Pty) Ltd			Version:			02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions
33	Imported and Exported	Vinceel	Liquid	1 x 20 mL	1 mL contains: The active substances are: Apis mellifica L D7 0,0125 mL; Atropa belladonna L D5 0,0125 mL; Calendula officinalis L D3 0,0125 mL; Echinacea purpurea (L.) Moench D5 0,0125 mL; Phytolacca americana L D3 0,0125 mL; Plantago major L D3 0,0125 mL; Salvia officinalis L D3 0,0125 mL; Vincetoxicum hirundinaria Medik D5 0,25 g. Preservative: Ethyl Alcohol 16,1 % m/m. The other ingredients are: Purified water, Glycerol.	The usual dose is: Not recommended for children below 6 years of age. In general, spray once (1 spray corresponds to 0,132-0,147 mL) 3 times a day. In acute conditions, spray once every 1/2 to every hour, up to 12 times a day.	D540660	Discipline specific	Homeopathy	Sublingual / Buccal	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in mouth and throat inflammations such as sore throats, gingivitis, pharyngitis, painful dentition, inflammation of the Eustachian tube and middle ear, tonsillitis.	At or below 25 °C
34	Imported and Exported	*[S3] Zeel T	Injectable (D5 only)	10 x 2 mL	Each 2 mL ampoule contains: The active substances are: Cartilago suis D6 2 mg; Funiculus umbilicalis suis D6 2 mg; Embryo suis D6 2 mg; Placenta suis D6 2 mg; Solanum dulcamara D3 10 mg; Symphytum officinale D6 10 mg; Nadidum D8 2 mg; Coenzymum A D8 2 mg; Sanguinaria canadensis D4 3 mg; Natrium diethyl oxalacetikum D8 2 mg; Acidum alpha-liponicum D8 2 mg; Toxicodendron quercifolium D2 10 mg; Arnica montana D4 200 mg; Sulfur D6 3,6 mg.	The usual dose is: In general, inject one ampoule twice weekly i.m., s.c., i.c., intra- or periarticularly, or, if necessary, paravertebrally. In the case of large joints, two ampoules on each occasion.	U5635	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in arthrosis (in particular gonarthrosis), polyarthrosis, spondylarthrosis, scapulohumeral periarthritis.	At or below 25 °C
35	Imported and Exported	Zeel T	Tablet	1 x 50	1 tablet cont.: The active substances are: Cartilago suis D4 0,3 mg; Funiculus umbilicalis suis D4 0,3 mg; Embryo suis D4 0,3 mg; Placenta suis D4 0,3 mg; Toxicodendron quercifolium D2 0,54 mg; Arnica montana D3 0,6 mg; Solanum dulcamara D2 0,15 mg; Symphytum officinale D8 0,15 mg; Sanguinaria canadensis D3 0,45 mg; Sulfur D6 0,54 mg; Nadidum D6 0,03 mg; Coenzymum A D6 0,03 mg; Natrium diethyl oxalacetikum D6 0,03 mg; Acidum alpha-liponicum D6 0,03 mg; Acidum silicicum D6 3 mg. The other ingredient is: Magnesium stearate. One tablet contains approx. 300 mg lactose.	The usual dose is: Adults: In general, 1 tablet to be dissolved in the mouth 3-5 times daily.	U5517	Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in arthrosis (in particular gonarthrosis), polyarthrosis, spondylarthrosis, scapulohumeral periarthritis.	At or below 25 °C
36	Imported and Exported	Nectadyn	Liquid	1 x 125 mL	Each 125 g contain: The active substances are: Baptista tinctoria D6 5,75 g; Drosera rotundi D4 6,0 g; Hydrastis canadensis D6 6,0 g; Ipecacuanha D4 6,0 g; Rumex crispus D 6,0 g; Sambucus nigra D4 6,0 g.	The usual dose is: ACUTE: Adults: 10 - 15 mL 3 - 5 times daily. Children (3 to 10 years old): 10 mL 3 - 5 times daily. Children (6 months to 3 years old): 5 mL 3 - 5 times daily. Babies (0 - 6 months): On the advice of a medical practitioner. CHRONIC: Adults: 5 mL 3 - 5 times daily. Children (3 - 10 years old): 2,5 mL 3-5 times daily. It is recommended to clean the enclosed measuring device after each use.	None	Discipline specific	Homeopathy	Oral	High	Relief from acute and chronic coughs of all origins, e.g. dry cough, expectorant cough, whooping cough, croup.	At or below 25 °C

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:		ModHomCo (Pty) Ltd			Version:			02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions
37	Imported and Exported	Euphorbium compositum	Liquid	1 x 20 mL	20 mL cont.: The active substances are: Euphorbium D4 0,04 mL; Pulsatilla pratensis D2 0,04 mL; Luffa operculata D2 0,04 mL; Mercurius bijodatus D8 0,04 mL; Mucosa nasalis suis (Sus scrofa ferus domesticus) D8 0,04 mL; Hepar sulfuris D10 0,04 mL; Argentum nitricum D10 0,04 mL; Sinusitis-Nosode D13 0,04 mL. The other ingredients are: Water, purified; Disodium phosphate dehydrate; Sodium chloride; Sodium dihydrogen phosphate dehydrate. Preservative: Benzalkonium chloride 0,01 %.	The usual dose is: Adults: 1-2 sprays into each nostril 3-5 times daily. Children 2-5 years: 1 spray into each nostril 3-4 times daily. Do not administer to children younger than 2 years.	U 5584 D540784	Discipline specific	Homeopathy	Nasal	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in rhinitis of varied origins (viral, bacterial, allergic); rhinitis sicca, rhinitis hyperplastica and atrophicans, for the auxiliary treatment of ozena; chronic sinusitis; to facilitate nasal respiration in hay fever.	At or below 25 °C
38	Imported and Exported	*[3] Traumeel LT ad us. vet.	Injectable (D5 only)	5 x 5 mL	Each 5 mL ampoule contains: Arnica Montana D4 500 mg; Atropa belladonna D4 500 mg; Calendula officinalis D4 500 mg; Aconitum napellus D4 300 mg; Bellis perennis D4 250 mg; Hypericum perforatum D4 150 mg; Echinacea D4 125 mg; Echinacea purpurea e planta tota D4 125 mg; Hamamelis virginiana D 4 50 mg; Achillea millefolium D5 500 mg; Matricaria recutita D5 500 mg; Hepar sulfuris D6 500 mg; Symphytum officinale D8 250 mg; Mercurius solubilis Hahnemanni D8 250 mg.	According to species, the single daily dosage is: Horse, cattle (over 500 kg): 10 ml once a day. Horse, cattle, swine (up to 500 kg) 5 ml once a day. Piglet: 2-3 ml once a day. Sheep, goat: 2 ml once a day. Large dog (over 25 kg): 3-4 ml once a day. Medium size dog (15-25 kg): 2 ml once a day. Small dog, cats (up to 15 kg): 1-2 ml once a day. Small pets: 0,5-1 ml once a day. If necessary, depending on the severity of the case, the indicated dosage is to be repeated after 24 hours. For long-term treatment of chronic affections or those with tendencies to recur, the indicated single dosage can be administered at 4-day intervals.		Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in injuries and their results, e.g. cerebral concussion, fractures, luxations, contusions, burns, postoperative and posttraumatic soft tissue swelling, edemas, hematoma. Inflammatory processes of different kinds and localisations, e.g. phlegmons, abscesses, inflammations of the anal sac, otitis, fluor albus (white vaginal discharge), dermatitis, conjunctivitis, mastitis. Inflammatory and degenerative processes accompanied by inflammations, especially at the connective and locomotorial system, e.g. arthritis, tendovaginitis, bursitis, galls of hockjoint.	At or below 25 °C
39	Imported and Exported	Traumeel T ad us. vet.	Tablet	1 x 100 1 x 250 1 x 500	1 tablet contains: Hypericum perforatum D2 3 mg; Aconitum napellus D3 30 mg; Chamomilla recutita D3 24 mg; Achillea millefolium D3 15 mg; Arnica montana D3 15 mg; Calendula officinalis D3 15 mg; Hamamelis virginiana D3 15 mg; Bellis perennis D3 6 mg; Echinacea D3 6 mg; Echinacea purpurea D3 6 mg; Atropa belladonna D4 75 mg; Mercurius solubilis Hahnemanni D8 30 mg; Hepar sulfuris D8 30 mg; Symphytum officinale D8 24 mg.	According to species, the daily dosage is: Horse, cattle: 2-3 times daily 6-10 tablets. Swine: 2-3 times daily 4 tablets. Sheep, goat: 2-3 times daily 2 tablets. Large dog: 2-3 times daily 3 tablets. Medium size dog: 2-3 times daily 2 tablets. Small dog, cat: 2-3 times daily 1 tablet. In cases with acute symptoms, initially the indicated dosage can be repeated after half to one hour during a period of 4 hours.		Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in injuries and their results, e.g. cerebral concussion, fractures, luxations, contusions, burns, postoperative and posttraumatic soft tissue swelling, edemas, hematoma. Inflammatory processes of different kinds and localisations, e.g. phlegmons, abscesses, inflammations of the anal sac, otitis, fluor albus (white vaginal discharge), dermatitis, conjunctivitis, mastitis. Inflammatory and degenerative processes accompanied by inflammations, especially at the connective and locomotorial system, e.g. arthritis, tendovaginitis, bursitis, galls of hockjoint.	At or below 25 °C
40	Imported and Exported	Traumeel Gel ad us. vet.	Gels	50 g 250 g	100 g contains: Calendula officinalis Ø 0.45 mg; Hamamelis virginiana Ø 0.45 mg; Echinacea Ø 0.15 g; Echinacea purpurea Ø 0.15 g; Matricaria recutita Ø 0.15 g; Bellis perennis Ø 0.1 g; Achillea millefolium Ø 0.09 g; Aconitum napellus D1 0.05 g; Atropa belladonna D1 0.05 g; Arnica montana D3 1.5 g; Symphytum officinale D4 0.1 g; Hypericum perforatum D6 0.09 g; Hepar sulfuris D6 0.025 g; Mercurius solubilis Hahnemanni D6 0.04 g.	The gel is completely massaged into the skin of the affected area several times a day (up to 4 times a day) until the disorder has disappeared.		Discipline specific	Homeopathy	Cutaneous	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in injuries and their results, e.g. cerebral concussion, fractures, luxations, contusions, burns, postoperative and posttraumatic soft tissue swelling, edemas, hematoma. Inflammatory processes of different kinds and localisations, e.g. phlegmons, abscesses, inflammations of the anal sac, otitis, fluor albus (white vaginal discharge), dermatitis, conjunctivitis, mastitis. Inflammatory and degenerative processes accompanied by inflammations, especially at the connective and locomotorial system, e.g. arthritis, tendovaginitis, bursitis, galls of hockjoint.	At or below 25 °C
41	Imported and Exported	*[3] Zeel ad us. vet.	Injectable (D5 only)	5 x 5 mL	Each 5mL ampoule contains: Rhus toxicodendron (Rhus toxicodendron L.) D2 25,0 mg; Arnica montana (Arnica montana L.) D3 50,0 mg; Solanum dulcamara (Solanum dulcamara L.) D3 25,0 mg; Sanguinaria canadensis (Sanguinaria canadensis L.) D4 7,5 mg; Symphytum officinale (Symphytum officinale L.) D6 25,0 mg; Sulfur D6 9,0 mg; Cartilago suis (Sus scrofa ferus domesticus) D6 5,0 mg; Embryo totalis suis (Sus scrofa ferus domesticus) D6 5,0 mg; Funiculus umbilicalis suis (Sus scrofa ferus domesticus) D6 5,0 mg; Placenta totalis suis (Sus scrofa ferus domesticus) D6 5,0 mg; Acidum thiocticum (thioctic acid) D8 5,0 mg; Coenzym A D8 5,0 mg; Nadidum (nadide) D8 5,0 mg; Natrium diethyloxalacetatum (sodium diethyloxalacetate) D8 5,0 mg.	According to species, the single daily dosage is: Horse, cattle (over 500 kg): 10 ml once a day. Horse, cattle, donkey (up to 500 kg): 5 ml once a day. Swine: 3 – 5 ml once a day. Sheep, goat: 3 – 4 ml once a day. Large dog (over 25 kg): 3 – 4 ml once a day. Medium size dog (15 – 25 kg): 2 ml once a day. Small dog, cat (up to 15 kg): 1 – 2 ml once a day. Small pets: 0,5 ml once a day. Administer single dose amount at 4-day intervals in cases of chronic disease for long-term therapy. In acute conditions the single dose may initially be repeated after 24 hours. For duration of treatment longer than 4 weeks a medicinal supervision is recommended.	D552033	Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in joint diseases, especially the chronic-degenerative type e.g. arthritis, especially at joints of hip, knee, shoulder and elbow, spondylitis, spondylarthrosis, disc prolapse and after a joint operation.	At or below 25 °C

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:		ModHomCo (Pty) Ltd			Version:			02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions
42	Imported and Exported	*[3] Ubichinon compositum ad us. vet.	Injectable (D5 only)	10 x 2,2 mL 100 x 2,2 mL	Each 2.2 mL ampoule contains: Colchicum autumnale D4; Conium maculatum D4; Hydrastis canadensis D4; Podophyllum peltatum D4; Vaccinium myrtillus D4; Acidum ascorbicum D6; Acidum sarcosolicum D6; Galium aparine D6; Natrium riboflavinum phosphoricum D6; Nicotinamidum D6; Pyridoxinum hydrochloricum D6; Thiaminum hydrochloricum D6; Acidum alpha-liponicum D8; Hydrochinonum D8; Manganum phosphoricum D8; Natrium diethyloxalacetum D8; Sulfur D8; Acidum acetylosalicylicum D10; Adenosinum thiphosphoricum D10; Antrachinonum D10; Coenzymum A D10; Histaminum D10; Magnesium glyconicum D10; Nadidum D10; Naphthochinonum D10; Para-Benzochinonum D10; Trichinoylum D10; Ubichinon D10; 22.0 mg each.	According to the species, the single daily dosage is: Large dog: 3-4 mL Medium dog: 2 mL Small dog, cat: 1-2 mL In the case of acute symptoms, the single dose indicated can initially be repeated after 12 hours, then continue with the standard dosage. For long-term treatment of chronic affections, the indicated single dosage can be administered at intervals of 1-4 days.		Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in chronic and degenerative diseases, organ degeneration, dysfunctions in cellular energy metabolism, improvement of oxygen utilisation, tumour diseases.	At or below 25 °C
43	Imported and Exported	*[3] Solidago compositum ad us. vet.	Injectable (D5 only)	10 x 2,2 mL 100 x 2,2 mL	Each 2.2 mL ampoule contains: Batisia tinctoria D4; Berberis vulgaris D4; Equisetum hyemale D4; Solidago virgaurea D4; Argentum nitricum D6; Capsicum annuum D6; Curpum sulfuricum D6; Lyttia vesicatoria D6; Orthosiphon aristatus D6; Pareira brava D6; SMilax D6; Terebinthina laricina D6; Apsinum D8; Bucco D8; Hydrargyrum bichloratum D8; Vesica urinaria suis D8; Hepar sulfuris D10; Natrium pyruvicum D10; Pyelon suis D10; Ureter suis D10; Acidum arsenicosum D12; 22 mg each.	According to species, the single daily dosage is: Large dog: 3-4 mL Medium dog: 2 mL Small dog, cat: 1-2 mL Puppy: 0.5-1 mL Small rodents: 0.5 mL Cage birds: 0.1-0.5 mL If necessary, depending on the severity of the case, the indicated dosage is to be repeated two times a day until two weeks. For treatment of disorders with tendencies to recur, chronic affections or long-term treatment, the indicated single dosage can be administered at 2 times a week intervals.		Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in acute and chronic diseases of the kidneys and urinary tract, such as cystitis, cystopyelitis, nephrolithiasis, hydronephrosis, nephrosis and nephrosclerosis, acute glomerulonephritis, the 1st stage of prostatic adenoma, incontinencia urinae, and for the stimulation of excretion from the kidneys, e.g. in the case of hyperhidrosis, sudoresis of the feet, eczema.	At or below 25 °C
44	Imported and Exported	Nurexan ad us. vet.	Tablet	1 x 50 1 x 100 1 x 500	1 tablet contains: Avena sativa DII. D2; Passiflora incarnata DII. D2; Yinzum Isolvalerianicum DII. D4; Coffea arabica DII. D12; 0,6 mg each.	According to species, the dosage is: Large dog (over 25 kg): 2-3 times daily the single dose of 3 tablets. Medium-sized dog (15-25 kg): 2-3 times daily the single dose of 2 tablets. Small dog (up to 15 kg), cat: 2-3 times daily the single dose of 1 tablet. For acute symptoms, initially administer the indicated single dose every half to full hour for up to 2 hours, then continue with the standard dose. For long-term treatment of chronic diseases, administer the single dose at intervals of 1 to 4 days.		Discipline specific	Homeopathy	Oral	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in nervous restlessness, sleep disturbances.	At or below 25 °C

SAHPRA LICENCE NO.		DL02-C93ACA	CM REF.	see Licence No.	Name of Licence Holder:		ModHomCo (Pty) Ltd						Version:		02-082023
No.	Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Sub-Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions		
45	Imported and Exported	*[3] Coenzyme ad us. vet	Injectable (DS only)	5 x 5 mL 50 x 5 mL	Each 5 mL ampoule contains: Beta vulgaris conditiva D4 0.05 mL; Acidum alphaliponicum D6 0.05 mL; Acidum ascorbicum D6 0.05 mL; Cysteinum D6 0.05 mL; Magnesium oroticum D6 0.05 mL; Manganum phosphoricum D6 0.05 mL; Natrium diethylolalacetum D6 0.05 mL; Natrium riboflavinum phosphoricum D6 0.05 mL; Nicotinamidum D6 0.05 mL; Pulsatilla pratensis D6 0.05 mL; Pyridoxinum hydrochloricum D6 0.05 mL; Thiaminum hydrochloricum D6 0.05 mL; Acidum alpha-ketoglutaricum D8 0.05 mL; Acidum cisaconiticum D8 0.05 mL; Acidum citricum D8 0.05 mL; Acidum DL malicum D8 0.05 mL; Acidum fumaricum D8 0.05 mL; Acidum succinicum D8 0.05 mL; Cerium oxalicum D8 0.05 mL; Coenzymum A D8 0.05 mL; Nadidum D8 0.05 mL; Natrium pyruvicum D8 0.05 mL; Adenosinum triphosphoricum D10 0.05 mL; Barium oxalsuccinicum D10 0.05 mL; Hepar sulfuris D10 0.05 mL; Sulfur D10 0.05 mL.	According to the species, the single daily dosage is: Horse, cattle, swine: 5 ml Piglet: 2-3 ml Sheep, goat: 2 ml Large dog: 3-4 ml Medium dog: 2 ml Small dog, cat: 1-2 ml Puppy: 0.5-1 ml Small pets: 0.5 ml In cases with acute symptoms, the indicated dosage can be repeated after a period of 24 hours. For treatment of disorders with tendencies to recur, chronic affections or long-term treatment, the indicated single dosage can be administered at 4-day intervals.		Discipline specific	Homeopathy	Injection	High	This medicine is prepared in accordance with homeopathic principles and is proposed for use in chronic and degenerative diseases, metabolic disorders, cachexia, enzyme blocks and dysfunctions in cellular metabolism, improvement of oxygen utilization, supportive in tumour diseases.	At or below 25 ° C		