

SAHPRA LICENCE NO		CM REF	DA02-922402	Name of Licence Holder	Abex Pharmaceutica						Version:	01-022021
o. Activity	Name of Medicine	Dosage Form	Pack size per unit - Quantity or Volume	Ingredients and Quantities	Dosage	Registration or Application No.	Category	Class or Discipline	Route of Administration	Risk	Intended Use	Storage Conditions
Imported and Exported	Sneezo	Tablet - Chewable	2 30 tablets	Zinc 0,75 mg (as Zinc bisglycinate 3,75 mg) per tablet; Quail egg powder 42,0 mg per tablet	SNEEZO can be used by adults and children from the age of 4 years. It is recommended that one to two tablets are chewed upon appearance of reactions to seasonal and environmental conditions. To be repeated as needed up to 5 times per day. SNEEZO can also be chewed 15 to 20 minutes before exposure to immune triggers.	Health Si		Multiple substance ormulation	Oral		Contributes to maintaining normal immune response to seasonal and environmental allergens	At or below 25 ° C
Imported and Exported	Tiger Balm Red	Other	19 g ointment	Natural camphor 11,0 % w/w; Levomenthol (menthol) 10,0 % w/w	Adults and children over 2 years of age: Rub gently on the affected parts of the skin as necessary, usually 2 or 3 times a day.	Health Si		Aultiple substance ormulation	Cutaneous	Low	Improves blood circulation in micro vessels     Decrease sensation of heavy legs*     Temporary relief of muscular cramps and rheumatic pain	At or below 25 * C
Imported and Exported	Tiger Balm White	Other	19 g ointment	Natural camphor 11,0 % w/w; Levomenthol (menthol) 8,0 % w/w	Adults and children over 2 years of age: Rub gently on the affected parts of the skin as necessary, usually 2 or 3 times a day.	Health Si	Supplement N	Multiple substance ormulation	Cutaneous	Low	Cooling sensation and/or soothing action for the skin     Temporary relief of mild ansal congestion     Temporary relief of chest congestion     Improves blood circulation in micro vessels     Temporary relief of the	At or below 25 * C
Imported and Exported	Pregula	Powder	30 powder sticks	Folic acid 200 µg per stick; Myo-inositol 2 g per stick	The recommended daily dose of PREGULA is 1 stick every 12 hours (2 sticks per day), to be poured directly into the mouth or to be dissolved in water. PREGULA should be taken 2 hours before or after a meal. PREGULA should be taken for at least 3 months.		Supplement N	Multiple substance ormulation	Oral	Low	To prepare and support pregnancy by helping to reduce the risk of neural tube defects when taken daily prior to becoming pregnant and during early pregnancy. Contributes to the maintenance of good health e.g. Maintains and supports a healthy female hormonal balance during the reproductive cycle Maintains, supports and regulates a healthy menstrual cycle.	At or below 25 ° C
Imported and Exported	Truvit C	Liquid	30 sachets	Sodium L ascorbate 1 000 mg per sachet	Adults and children over 9 years of age: Take 1 sachet once daily orally.	Health Si	Supplement V	/itamins	Oral	Low	Contributes to cell protection from free radical damage Contributes to normal collagen formation for the normal function of bones, teeth, skin and cardiage Contributes to the normal function of the immune system Contributes to the reduction of tiredness and fatigue A factor in the maintenance of good health	At or below 25 ° C
Imported and Exported	Truvit D 1000 IU	Liquid	30 sachets	1 000 IU cholecalciferol, equivalent to 25 micrograms vitamin D3 per sachet	Adults and children over 1 year of age: Take 1 sachet once daily orally	Health Si	Supplement V	/itamins	Oral	Low	Helps in the absorption and use of calcium and phosphorous.     Contributes to the development and maintenance of strong bones and teeth.     calcium intake, when combined with sufficient vitamin D, a healthy diet and regular exercise, may reduce the risk of developing osteoporosis.	At or below 25 ° C
Imported and Exported	Truvit D 400 IU	Liquid	30 sachets	400 IU cholecalciferol, equivalent to 10 micrograms vitamin D3 per sachet	Adults and children over 1 year of age: Take 1 sachet once daily orally	Health Si	Supplement V	/itamins	Oral			At or below 25 * C